

The Facts about Head Lice

Further copies of this leaflet are available FREE from:

Publications Centre

The National Assembly for Wales

Cathays Park

Cardiff

CF10 3NQ

Email: assembly-publications@wales.gsi.gov.uk

@Crown copyright 2008

CMK-22-02-224

G/755/07-08

Fax: (029) 2082 5239

What are head lice?

- Head lice are small insects (no larger than a sesame seed when fully grown) with moving legs. They live very close to the scalp.
- Nits are not the same as lice. They are the empty dead egg cases which stick to the hair, and look like specks of dandruff.
- You only have head lice if you find a moving louse (not a nit). Your head does not have to be itchy to have head lice.

How do you get head lice?

- Because of children's play activity and close contact, head lice are usually found on children, but can also spread to adults.
- Head lice infections are caught from close family and friends in the home and community, as well as at school.
- Head lice live in all types of hair – dirty or clean, short or long.
- Head lice walk from one head to another, when heads are close together - they do not fly, jump or swim.

[These notes are intended for families and may be photocopied for circulation](#)

How can you prevent head lice?

- Regular brushing and combing of the hair will help to spot lice early.
- The best way to stop the spread of head lice is for families to check their heads by regular wet combing using a detector comb.
- Human head lice are not found on animals or household pets and are not transmitted from pets to humans.

What are the side effects of head lice?

The reaction of individuals to louse bites can vary enormously from person to person. When you have head lice for the first time, you will probably not even realise they are there. However, after a short time (up to a few weeks), you start to itch, and react to the bites. Once you have had this initial reaction, you will experience itching and redness within 24-48 hours, each time you become re-infested. However, some individuals never exhibit symptoms. Itching from large numbers of bites may make the infested individual feel tired, irritable and feverish, hence the term 'feeling lousy'.

How do you detect head lice?

YOU WILL NEED: A DETECTOR COMB

[\(These can be bought from your pharmacist - ask for help if there are none on display\).](#)

How do you treat head lice?

DO NOT TREAT UNLESS YOU ARE SURE YOU HAVE FOUND A MOVING LOUSE.

1. Check and treat all infected members of the family.
2. Ask your GP or Pharmacist which head lice treatment to use. Do not use head lice shampoo. The Pharmacist can advise on any evidence of resistance to treatment and therefore the most appropriate treatment to purchase. Also, the Pharmacist can advise on treatments suitable for people with asthma and other respiratory conditions.
3. Always read and follow the instructions in the treatment pack.
4. Apply the treatment carefully.

The usual methods are: Lotions and Liquids

In a well ventilated room (open a window), and away from all sources of heat (fi res, cigarettes, hairdryers, lighters or matches).

- i. Apply the lotion or liquid to dry hair
- ii. Part the hair near the top of the head, put a few drops of the lotion or liquid on to the scalp and rub in. Part the hair again a bit further down the scalp and rub in some of the lotion or liquid. Do this again and again until the whole scalp is wet.

If you are treating long hair, ensure that you cover the hair down to where the ponytail band would be, UNLESS you are using Dimeticone (Hedrin) which should fully cover the hair. Ensure that you use enough lotion - do not dilute head lice treatments.

Take care not to get the lotion or liquid in the eyes or on the face.

- iii. Let the lotion or liquid dry on the hair naturally. Keep well away from naked flames, cigarettes or other lighted objects.

Do NOT use a hair dryer.

You should use at least one small bottle of lotion or liquid per head, more if the hair is thick.

How can you carry out contact tracing?

Head lice are likely to have been picked up the lice from a close contact, usually a family member or close friend, who probably doesn't know they have lice. There should be no reason to feel embarrassed and it is best to inform likely contacts as soon as possible to prevent further spread.

What if the problem won't go away?

Did You ...

- use enough lotion/liquid?
- apply it correctly?
- let it dry naturally?
- leave it on for the correct time?
- use a second bottle 7 days after the first?
- check all your family and friends?
- check adults as well as children?
- treat all infected contacts at the same time?

Why do we no longer have a nit nurse?

It has been recognised that routine inspections at school by a school nurse are not only ineffective at tackling head lice in the long term, but can lead to a false sense of security. The responsibility for frequent checks must lie with the individual parent/s (who is best placed to detect infestation).

In order to encourage this method the Parent Teacher Associations of some schools organise family health promotion sessions with health professionals. Many schools will participate in National Bug Busting Days

(31 January, 15 June and 31 October) which aim to ensure children and parents know how to detect and treat head lice and stop their circulation.

Where can you go for further help and advice?

Contact the local GP, Pharmacist or Health Protection Nurse.

Do those infected with head lice need to stay indoors until the head lice are gone?

There is no need to keep those suffering from head lice indoors, or children from attending school.

Should a child be excluded from school if they have head lice?

No. Exclusion is a disciplinary measure that should be used in exceptional circumstances and as a last resort and should not be used to manage head lice infection.

Should other parents be notified if a child is infected with head lice in their child's school?

Despite popular belief, letters notifying other parents of cases of head lice do not have any effect in stopping the spread of lice amongst children. However, letters of this kind often have the effect of causing itching and anxiety as a purely psychological response.

Are lice shampoos potentially hazardous?

There are possible side effects/hazards from using lice shampoos e.g. those with respiratory conditions such as asthma, epilepsy, pregnant women etc. Advice should be sought from your local GP or Pharmacist who will be able to advise further.

Do I need to spray my bedding and furniture?

Head lice are human parasites and require human blood to survive.

They are not environmental pests so pesticide sprays for furniture and bedding are unnecessary and a serious risk to health. Vacuuming is the safest and best way to remove lice or fallen hairs with attached nits from upholstered furniture, rugs, stuffed animals and cars.

Do pets get head lice?

Do head lice spread diseases?

No, head lice are not known to transmit any disease agents. However, if the skin is broken by repeated scratching, other infections can be introduced by organisms under the nails, but this is extremely uncommon.

References

Public Health Medicine Environmental Group, (2008), Head Lice: Evidence-Based Guidelines Based on the Stafford Report 2008 Update. Illustrations courtesy of the health charity www.chc.org/bugbusting and www.headlice.org.uk